## Wisconsin Nightjar Survey, 2008-2011

	WHIP-POOR-WILL									
	2008-1	2008-2	2009-1	2009-2	2010-1	2010-2	2011-1	2011-2		
# routes surveyed	71		72	29	62	24	53			
# birds detected	169		167	72	141	36	155			
# birds per route	2.38		2.32	2.48	2.27	1.50	2.92			
# routes w/ target	23		24	9	23	7	17			
# birds per route w/ target	7.35		6.96	8.00	6.13	5.14	9.12			
# routes = 0 birds	48		48	20	39	17	36			
# routes = 1-5 birds	16		12	2	13	5	8			
# routes = 6-10 birds	2		8	5	5	1	2			
# routes > 10 birds	5		4	2	5	1	7			
Top 3 routes (# birds)	#11 (33) #29 (26) #137 (26)	No surveys conducted	#11 (25) #137 (22) #28, 43 (15)	#28 (13) #137 (13) #43 (10)	#28 (19) #137 (16) #29 (16)	#137 (15) #41 (8) #22, 38 (4)	#11 (24) #28 (24) #137 (24)	No surveys conducted		

	COMMON NIGHTHAWK									
	2008-1	2008-2	2009-1	2009-2	2010-1	2010-2	2011-1	2011-2		
# routes surveyed	71		72	29	62	24	53			
# birds detected	7		42	4	14	0	6			
# birds per route	0.10		0.58	0.14	0.23	0.00	0.11			
# routes w/ target	6		15	2	10	0	4			
# birds per route w/ target	1.17		2.80	2.00	1.4	0	1.5			
# routes = 0 birds	65		57	27	52	24	49			
# routes = 1-5 birds	6		14	2	10	0	4			
# routes = 6-10 birds	0		0	0	0	0	0			
# routes > 10 birds	0		1	0	0	0	0			
Top 3 routes (# birds)	#33 (2)	No surveys conducted	#69 (20) #26 (4) #301 (4)	#11 (3) #303 (1)	#40 (3) #56 (2) #118 (2)	n/a	#30 (2) #65 (2)	No surveys conducted		